

Maxillofacial Injuries Associated With Domestic Violence

Bach T. Le, DDS, MD, Eric J. Dierks, DMD, MD,†
Brett A. Ueeck, DMD,‡ Louis D. Homer, MD, PhD,§
and Bryce F. Potter, DMD, MD||*

Purpose: The purpose of this study was to report the incidence, causes, and patterns of maxillofacial injury associated with domestic violence.

Patients and Methods: A retrospective review of patients treated for domestic violence injuries at an inner-city hospital over a 5-year period was done, and data were collected on type and location of injury, mechanism of injury, alcohol involvement, and treatment.

Results: The sample consisted of 236 emergency room admissions. The majority (81%) of victims presented with maxillofacial injuries. The fist was a favorite means for assaults (67%). The middle third of the face was most commonly involved (69%). Soft tissue injuries were the most common type of injury (61%). Facial fractures were present in 30% of victims. The average number of mandible fractures per patient was 1.32. The majority of facial fractures (40%) were nasal fractures. Left-sided facial injuries were more common than right sided.

Conclusions: These data confirm that most victims of domestic violence sustain maxillofacial injuries. Midface injuries predominate. The preponderance of facial injuries makes it very likely that oral and maxillofacial surgeons will be involved in the care of these patients.

© 2001 American Association of Oral and Maxillofacial Surgeons

Domestic violence afflicts millions of people each year. A woman has a greater than 1 in 5 chance of being injured during such violence.¹ In the United States, 2 to 4 million women are assaulted by their intimate partners annually, and domestic violence is currently the most common cause of nonfatal injury in women.² Currently, 1 in 3 homicides in the United States are a direct result from domestic violence.³

Aside from the knowledge that most injuries resulting from domestic violence involve the face, there is little information about facial injury patterns that oc-

cur. This article attempts to identify the specific patterns of maxillofacial injuries that are commonly seen in such victims.

Patients and Methods

The records of 236 patients treated for domestic violence injuries between the beginning of January 1992 and the end of December 1996 at an inner-city level I trauma hospital (Legacy Emanuel Hospital, Portland, OR) were retrospectively reviewed. Using the Emanuel ER Registry, information was requested on all women who gave a positive history of being intentionally injured by their spouse or sexual partner during this period. Data were collected on type and location of injury, mechanism of injury, alcohol involvement, and treatment.

Injuries were recorded according to anatomic location as head, maxillofacial, neck, breast, chest, abdomen, back, buttocks, and extremities. The injuries were also classified as contusions and abrasions, lacerations, and fractures and dislocations. Facial injuries were classified according to location, type, lateralization (left vs right), and facial third. The data were analyzed and tested for statistical significance using descriptive statistics and the chi-square test.

*Formerly, Chief Resident, Department of Oral and Maxillofacial Surgery, Oregon Health Sciences University, Portland, OR; Currently, Assistant Professor, LAC-USC Medical Center, Los Angeles, CA.

†Vice Chairman, Department of Oral and Maxillofacial Surgery, Oregon Health Sciences University, Portland, OR.

‡Resident, Department of Oral and Maxillofacial Surgery, Oregon Health Sciences University, Portland, OR.

§Medical Director of Clinical Investigations & Biomedical Research, Legacy Health Systems, Portland, OR.

||Clinical Professor, Department of Oral and Maxillofacial Surgery, Oregon Health Sciences University, Portland, OR.

Address correspondence and reprint requests to Dr Le: 273 East Glenarm St, #5, Pasadena, CA 91106; e-mail: Leb97201@yahoo.com

© 2001 American Association of Oral and Maxillofacial Surgeons

0278-2391/01/5911-0004\$35.00/0

doi:10.1053/joms.2001.27490




FIGURE 1. Distribution of contusions and abrasions.




FIGURE 3. Distribution of fractures and dislocations.

Results

The results are based on the records of 236 consecutive emergency room admissions for domestic violence. Mean patient age was 31.4 years, with a range between 15 and 71 years. All cases were females. One hundred fifty-five (66%) patients had reported previous abuse. Thirty-four patients (14%) had injuries severe enough to require admission to the hospital. One patient died as a result of her injuries.

The majority (78%) of victims were single, separated, or divorced. One hundred and seventy-nine (76%) patients were unemployed. Alcohol was involved in 33% of the cases. One hundred and forty-two (60%) patients had a drug abuse history. Nine patients were pregnant at the time of assault.

The 236 women had a total of 257 contusions and abrasions, 70 lacerations, and 93 fractures and dislocations (Figs 1 to 3). Most injuries were located on the face. Eighty-one percent of victims presented with maxillofacial injuries. Fifty percent of the study population had an isolated maxillofacial injury as the only presenting trauma. The remaining cases (31%) had multiple presenting injuries, whereas only 14% had an isolated nonmaxillofacial injury (Fig 4).

Table 1 shows the location of the facial injuries encountered. Although there were 236 patients, the numbers add up to more than 236 because some victims had more than one injury recorded. Accordingly, the percentages total more than 100%. A “none” category was created to account for patients without facial injuries.




FIGURE 2. Distribution of lacerations.

Soft tissue injuries in the form of contusions of the maxillofacial region were the most common types of injury (61%). Forty patients (17%) had facial lacerations serious enough to require repair. The middle third of the face was most commonly involved (69%), followed by upper third (13%), and lower third (19%). Patients with facial injury locations that were not clearly specified were labeled “not specified.”

Seventy patients (30%) sustained 85 facial fractures. Most facial fractures were nasal fractures (40%). One fracture (1.2%) involved the upper face only, 57 fractures (67%) involved the middle face only, and 27 fractures (32%) involved the lower face only. The one upper fracture was isolated. Midface fractures were isolated in 39 of 57 instances, but were present as 2 fractures in 8 cases, and as 3 fractures in 1 case (Table 2). The majority of middle third fractures sustained were also nasal fractures (68%).




FIGURE 4. Distribution of injuries.

Table 1. DISTRIBUTION OF FACIAL INJURIES

| | Fracture Upper Face | Fracture Midface | Fracture Lower Face | L/C Upper Face | L/C Midface | L/C Lower Face | None /Not Specified |
|------------|------------------------|---------------------|------------------------|-------------------|----------------|-------------------|------------------------|
| No. | 1.00 | 57.00 | 27.00 | 30.00 | 105.00 | 17.00 | 46.00 |
| Percentage | 0.42 | 24.15 | 11.44 | 12.71 | 44.49 | 7.20 | 19.49 |

Abbreviations: L, laceration; C, contusion.

The mandible fractures were a single fracture in 17 cases; there were double fractures in 2 cases and 4 fractures in 1 case. The average number of fractures per patient was 1.32. It appears that multiple fractures were most often located in the midface (Table 2). A likelihood ratio chi-square test of the hypothesis that multiple fractures were equally common in each region could not be rejected (chi-square = 2, 2 degrees of freedom, $P = .3$). When the question was asked whether a fracture was just as likely to be located in one region as in another, each with a probability of 1 in 3, the likelihood ratio test for this null hypothesis has a chi-square statistic of 77 with 2 degrees of freedom, $P < .000001$, indicating that there is a predilection for the midface over the upper facial region.

The fractures were further classified as midline (34), right-sided (10), or left-sided (43) (Table 3). Left-sided facial fractures were more common than right-sided fractures. Among the lateralized fractures, 43 of 53 (81%) were on the left, with an upper 95% confidence limit of 89% and a lower 95% confidence limit of 70%. The null hypothesis that right-sided and left-sided fractures were equally likely was easily rejected using the binomial distribution ($P < .0001$). Figure 5 shows the distribution of facial fractures among the 70 patients who sustained facial fractures.

Table 2. DISTRIBUTION AND NUMBER OF FACIAL FRACTURES

| Areas Fractured | No. of Fractures |
|----------------------------|------------------|
| Upper face only | 1 |
| Middle face (n = 57) | |
| Isolated midface | 39 |
| 2 fractures of midface | 8 |
| 3 fractures of midface | 1 |
| Lower face (n = 25) | |
| Isolated mandible | 17 |
| 2 fractures of mandible | 2 |
| 3 fractures of mandible | 0 |
| 4 fractures of mandible | 4 |
| Upper face and middle face | 0 |
| Middle face and lower face | 3 |
| Upper face and lower face | 0 |
| All regions | 0 |

Because some patients had more than 1 fracture, the total percentage totaled more than 100%.

MECHANISMS OF INJURY

Injuries were inflicted by either blunt or penetrating forces or a combination of both. Multiple sources of trauma accounted for the number of injuries being greater than the 236 victims. The fist was a favorite means of assault (67%) in this study population. A weapon was used to inflict injuries in 36 patients (15%). Most of the weapons were blunt objects, such as bottles, sticks, and pipes. A knife was used in 8 cases. Twenty-two women were kicked, and 8 women were choked. One woman suffered a gunshot wound. Figure 6 lists the frequency of the different mechanisms of injury in this study population.

Table 4 shows the cross tabulation of mechanism of injury with location and type of injury. Again, this shows the fist as the preferred mechanism of injury and the midface as the most frequent target.* No statistical evaluation of whether the coincidence of the 2 factors was greater than might be expected by chance was done. The usual tests for chance assortment require that the categories being examined are mutually exclusive. It should be noted that with 45% in the LCM category and 68% in the fist category, a chance assortment would give 45×68 (31%), which is not far from the 35% actually recorded.

Alcohol had been consumed by the victims in 77 cases (33%) at the time of assault. Alcohol use by the assailants was not known. When the use of alcohol was not documented it was assumed not to have been a factor. The association of alcohol with the mechanism of injury is shown in Table 5. The percentage of cases in which alcohol was a factor in these categories was not statistically different from the overall rate of 33%. Table 6 shows a similar association between injury categories and alcohol as a factor. Again no

* Because a visit may appear more than once in a row or column, the total number of entries exceeds the 236 cases whose records were reviewed. Also, some visits had no entries in these categories, meaning no maxillofacial injuries were recorded, and this is noted in the rows and columns labeled "NONE."

Table 3. MAXILLOFACIAL SKELETAL FRACTURE DISTRIBUTION

| Fracture Types | Subtype | Right Side | Left Side | Frequency Fractures |
|-------------------|-------------------------|------------|-----------|---------------------|
| Temporal* (n = 2) | | 1 | 1 | 2 |
| Orbit (n = 16) | Rims | | 1 | 1 |
| | Blow out | 3 | 12 | 15 |
| Nasal (n = 28) | | | | 28 (midline) |
| Maxilla (n = 0) | Le Fort I | | | 0 |
| | Le Fort II | | | 0 |
| | Le Fort III | | | 0 |
| Zygoma (n = 14) | Body/complex | | 12 | 12 |
| | Arch | 1 | 2 | 3 |
| Mandible (n = 25) | Condyle | 4 | 4 | 8 |
| | Coronoid | | | 0 |
| | Ramus | | 1 | 1 |
| | Body | | 3 | 3 |
| | Angle | 1 | 6 | 7 |
| | Symphysis/parasymphysis | | | 4 (midline) |
| | Alveolar | | | 2 (midline) |

*Two women sustained temporal bone fractures from blunt injuries with weapons.

percentages were very different from the overall percentage of 33%.

Of the 77 cases in which alcohol use was involved, 62 (81%) had a total of 78 injuries related to the maxillofacial region. Twenty (30%) of these patients had a total of 25 facial fractures, and 15 (19%) patients had no maxillofacial injury.

Sixty-six percent of the 236 patients required only minor treatment and were released. The remaining 34% received treatment that required a procedure. Thirty-one patients (14%) had injuries severe enough to require admission to the hospital.

Eighty-one percent of the patients had maxillofacial injuries as a component of their injury. Those patients with more serious maxillofacial injuries were referred to the oral and maxillofacial surgery

service for further evaluation and treatment. Of the midface fractures, 76% did not require treatment. When a procedure was undertaken, 60% were open procedures, whereas 40% were treated closed. Of the mandible fractures, 18.4% were not treated. When the mandible fractures were treated, 71.4% were treated by an open procedure, and 28.6% were treated as a closed procedure. Of the 70 patients who sustained facial fractures, 24.2% were treated in the operating room, whether it was an open or closed procedure.

Most untreated facial fractures were nasal fractures that were minimally displaced. Most patients were followed on an outpatient basis and may require surgery at a later time. These data do not reflect the follow-up treatment of these patients.




FIGURE 5. Distribution of facial fractures.




FIGURE 6. Mechanism of injury during abuse.

Most patients (86%) were discharged to home after the emergency department rendered medical services. Six patients left the hospital against medical advice before treatment could be rendered. The 34 patients admitted to the hospital had a length of stay (LOS) totaling 126 days (average LOS of 3.7 days per patient; 1 patient had a 50-day LOS). In most of these patients (85%), the primary reason for admission was related to the treatment of oral and maxillofacial surgery injuries, mainly maxillofacial fractures (zygoma and mandible fractures). Most patients (68%) were admitted to the oral and maxillofacial surgery service, whereas 8 were admitted to the trauma service, 2 to the obstetrics-gynecology service, and 1 to the general medicine service. One patient died from her injuries, and 1 patient became a quadriplegic as a result of her injury.

Discussion

This is a retrospective study of 236 consecutive emergency room visits of patients who sustained injuries as a result of domestic violence. The patients included in this study were obtained from the LEHHC Patient Registry. Data for this registry, started in 1983, are collected at multiple points of entry. The data are entered and E-coded based on the nature of the disease, mechanism of injury, trauma, etc. Only those

Table 5. ASSOCIATION OF ALCOHOL WITH MECHANISM OF INJURY

| | Sexual Assault | Fist | Bite | Weapon | Kick | Choke | None/Not Specified |
|---------------------|----------------|-------|------|--------|------|-------|--------------------|
| No. Alcohol present | 3 | 160 | 3 | 36 | 22 | 8 | 30 |
| | 0% | 31.3% | 0% | 44% | 32% | 13% | 30% |

patients designated through an E code as injured by interpersonal violence by a sexual partner were included.

The limitations of a retrospective study include recall bias and misclassification of study variables. Furthermore, the assessment based on chart review may underestimate the problem of drug and alcohol abuse, because these variables could only be included if they were clearly documented. Fortunately, the LEHHC patient registry has detailed and accurate records of most of the variables that were scrutinized. The emergency room physicians or registered nurses complete data input, rather than it being entered by a coding technician who has never seen the patient. In addition, the data entered is double-checked by computer verification software (Collector Digital Innovation Inc, Belair, MD), which completes up to 100 routine data checks looking for contradictions. The registry coordinator is responsible for the database and completes the final verification in this process.

This study showed that most women victims of domestic violence sustain a high number of maxillofacial injuries, supporting the study hypothesis. The upper limb was also a common site of injury. This may reflect the tendency of victims to defend themselves with their hands during the assault.

The pattern of facial injury was similar to that in other studies on assault victims.^{4,7} Some explanations for this include preference and accessibility of the face as a target for assailants. The proportion of fractures of the middle face, mainly nasal fractures, was very high. This was because of the prominent projection of the nose from the face and to the fact that it

Table 4. CROSS-TABULATION OF MECHANISM OF INJURY VERSUS LOCATION AND TYPE OF INJURY

| Mechanism of Injury | Fracture Upper Face | Fracture Midface | Fracture Lower Face | C/L Upper Face | C/L Midface | C/L Lower Face | None/Not Specified |
|---------------------|---------------------|------------------|---------------------|----------------|-------------|----------------|--------------------|
| Sexual assault | 0 | 0 | .42 | 0 | .4 | 0.4 | .85 |
| Fist | 0 | 20 | 9.3 | 6.8 | 35.2 | 6 | 5.9 |
| Bite | 0 | 0 | 0 | 0 | .85 | 0 | .42 |
| Foreign object | .4 | 3.4 | .85 | 3.4 | 6.4 | .4 | 4.2 |
| Kicked | 0 | 1.3 | 0 | .85 | 4.7 | .85 | 3 |
| Choked | 0 | 0 | 0 | .85 | 1.7 | 0.4 | .85 |
| Other/not specified | 0 | 0 | .85 | 2.1 | 2.5 | .85 | 5.9 |

Abbreviations: L, laceration, C, contusion.

Table 6. NATURE OF FACIAL INJURY AND ALCOHOL

| | Facial Fractures | Facial Lacerations and Contusions | None/Not Specified |
|-----------------|------------------|-----------------------------------|--------------------|
| No. | 85 | 152 | 46.00 |
| Alcohol present | 33% | 38% | 37% |

takes less force to fracture the nasal bones than other facial bones. The left zygoma was the second most commonly fractured facial bone. This probably reflects the fact that the most common mechanism of injury was the fist and that more than 90% of the population is right-handed.⁸ Furthermore, Grinker and Saks⁹ suggest that hemispheric cerebral dominance leads the victim to turn to the right in a reflexive manner to avoid being punched, thus presenting the left side of the face to the assailant.

The average number of facial fractures per patient was 1.21. Larger series of facial fractures have found higher numbers of fractures per patient, ranging from 1.6 to 1.8.^{10,11} The lower average number of fractures per patient could be explained by the fact that the mechanism of injury in this study population was predominantly assault, primarily using the fist. Other studies reporting facial fracture patterns include multiple etiologies, such as motor vehicle accidents, that tend to cause a greater number of fractures per patient because of a higher force of impact. In this series, midface fractures were more common than mandible fractures, whether nasal fractures were counted or excluded. Fractures of the condylar process showed the highest incidence of all mandibular fractures. This was in contrast to other series in which a predominance of fractures of the body of the mandible was reported. This difference may have been caused by the small number of mandible fractures in this study.

The use of alcohol and/or illicit drugs has been a controversial topic in domestic violence. Some clinicians believe that they are causative in both violent acts and victimization, whereas others believe that alcohol and drug abuse by the victims may be a result of the violence.¹²⁻¹⁶ There is no question that there is an association. Alcohol was found to be associated with at least one third of the assaults and more than half of our study group had a history of illicit drug use. Again, in a retrospective study, this is an under-assessment. It is suspected that many more of the patients were under the influence of alcohol at the time of abuse. In a prospective study of patients with facial fractures, McDade et al¹⁷ found that half of them showed some degree of alcohol dependence and that 60% of them sustained fractures as a result of an assault.

Clearly there is evidence of increased mortality and morbidity in patients who abuse alcohol and present with facial injuries.¹⁸ It is important to note this association with alcohol and drug abuse because it has significant implications in the immediate management of the patient.

Although it was thought that the patterns of injury would be more severe with the introduction of alcohol, no relationship was noted. These findings are different from those of Hutchison et al¹⁹ who found in their prospective study of a large number of patients with facial injuries that alcohol consumption was associated with more serious facial injuries.

Domestic violence is a source of considerable morbidity. The spectrum of injuries in our patients was broad. Most patients were treated for minor contusions and abrasions. Many victims required repair of lacerations (primarily on the face) and casting of broken and dislocated bones. Others required admission to the hospital for treatment of more serious injuries. When considering the fact that many of these patients repeatedly access the healthcare system before they are finally identified, it is important to recognize such victims early, before the escalating violence leads to more severe injuries or death. Asking directly about violence as a cause or contributing factor to a woman's injuries may save time, money, and the patient.

Although family physicians, obstetricians, and emergency physicians are sometimes considered the most likely to see battered women, oral and maxillofacial surgeons are also likely to see these patients because of the severe facial injuries. Of the patients who required hospital admissions, the majority was admitted to the oral and maxillofacial surgery service for treatment of their facial injuries.

The preponderance of facial injuries in this study makes it very likely that the oral and maxillofacial surgeon will be involved in the care of these patients. It is therefore important that we as oral and maxillofacial surgeons be cognizant to the broad social implication of domestic violence. Appropriate referrals are mandatory in the overall care of these patients. The proper identification and referral of victims will be greatly expedited if clinicians become more aware of the prevalence of domestic violence and more alert to its risk factors.

References

1. Wilt S, Olson S: Prevalence of domestic violence in the United States. *J Am Med Womens Assoc* 51:77, 1996
2. Grisso JA, Wishner AR, Schwartz DF, et al: A population-based study of injuries in inner-city women. *Am J Epidemiol* 134:59, 1991
3. Kellermann AL, Mercy JA: Men, women, and murder: Gender-specific differences in rates of fatal violence and victimization. *J Trauma* 33:1, 1992

4. Andersson L, Hultin M, Nordenram A, et al: Jaw fractures in the county of Stockholm. *Int J Oral Surg* 13:194, 1984
5. Hill CM, Crosher RF, Carroll MJ, et al: Facial fractures: The results of a prospective four-year study. *J Maxillofac Surg* 12:267, 1984
6. McLennan WD: Fractures of the malar (zygomatic) bone. *J R Coll Surg Edinb* 22:187, 1977
7. Voss R: Changing etiologic pattern of jaw fractures, *in* Jacobs JR (ed): *Maxillofacial Trauma: An International Perspective*. New York, NY, Praeger, 1983, p 126
8. Dejong NR: *The Neurologic Examination* (ed 2). New York, NY, Pitman Medical Publishing Co, 1958
9. Grinker RR, Saks AL: The cerebrum-cerebral hemispheres, *in* *Neurology* (ed 6). Springfield, IL, Charles C. Thomas, 1984, p 142
10. Van Hoof RF, Merkx CA, Stekelenburg C: The different patterns of fractures of the facial skeleton in four European countries. *Int J Oral Surg* 6:3, 1977
11. Hagan EH, Huelke DF: An analysis of 319 case reports of mandibular fractures. *J Oral Surg* 9:23, 1961
12. Lindman R, von der Pahlen B, Ost B, et al: Serum testosterone, cortisol, glucose, and ethanol in males arrested for spouse abuse. *Aggressive Behav* 18:393, 1992
13. Hotaling GT, Sugarman DB: An analysis of risk markers in husband to wife violence: The current state of knowledge. *Violence Vict* 1:101, 1986
14. Gustafson R: Alcohol and aggression: A replication study controlling for potential confounding variables. *Aggressive Behav* 18:21, 1992
15. Gantner AB, Taylor SP: Human physical aggression as a function of alcohol and threat of harm. *Aggressive Behav* 18:29, 1992
16. Cloninger CR: Neurogenetic adaptive mechanisms in alcoholism. *Science* 236:410, 1987
17. McDade DM, McNicol RD, Ward-Booth P, et al: The aetiology of maxillo-facial injuries, with special reference to the abuse of alcohol. *Int J Oral Surg* 11:152, 1982
18. McInnes GP, Young RE, Areny BS: Cardiac arrest following chlormethiazole infusion in chronic alcoholics. *Postgrad Med J* 50:724, 1980
19. Hutchison IL, Magennis P, Shepherd JP, et al: The BAOMS United Kingdom Survey of Facial Injuries Part 1: Aetiology and the association with alcohol consumption. *Br J Oral Maxillofac Surg* 36:3, 1986